

**NET REVISION**

**1<sup>ST</sup> SECONDARY**

**PHYSICS**

## Part (1) Questions

### 1) Complete the following statements:

- 1- When a force acts on a moving body in the same direction, its speed ..... Without changing .....
- 2- When a force acts on a moving body in an opposite direction of its motion, its speed ..... without changing .....
- 3- If a body moves in a circular path, its velocity changes in .....
- 4- If the radius of curvature of the path of a body moving in a circular path increases four times , so the centripetal force required to make the speed of the body constant decreases to .....
- 5- From life applications of centripetal force are ....., ..... and .....
- 6- The centripetal force acting on a satellite of mass (m) rotates around Earth with velocity (v) in constant orbit of a distance (r) from Earth's center equals .....
- 7- The universal gravitational constant is determined using the relation  $G = \dots\dots\dots$
- 8- The ratio between the universal gravitational constant on Earth's surface to that on Moon's surface is ..... unity.
- 9- If the distance between the centers of two bodies is doubled the force of attraction between them becomes its .....value.
- 10- ..... is the unit for measurement the universal gravitational constant.

## 2) Q2: What's meant by?

- 1- Uniform circular motion
- 2- Centripetal force.
- 3- Tangential velocity
- 4- Reaction force
- 5- Centripetal acceleration
- 6- Periodic time
- 7- Universal gravitational law
- 8- Gravitational field
- 9- Gravitational field intensity
- 10- Satellite
- 11- Orbital velocity
- 12- Astronomical moon

## 3) Q3: Give reason:


- 1- Although the object moves in a uniform circular motion and is affected by an acceleration, it has a linear constant velocity.
- 2- When a car turns in a circular path, it conserves moving in the curved path.
- 3- Although the body moving in a circular path is affected by centripetal force, it doesn't go to center of the circle.
- 4- Racer leans his body with the motorbike towards the center of the circular path at the curved part of the path.
- 5- The driver must decrease his velocity in the curved path.
- 6- The attraction force between two masses increases as they move nearer to each other.
- 7- The orbital velocity keeps the satellite at the same height.
- 8- The importance of satellites.

4) **Q4: What do we mean when we say?**

- 1- The centripetal force acting on an object is 100N.
- 2- The centripetal acceleration of an object is  $40 \text{ m/s}^2$ .
- 3- The orbital velocity of a satellite =  $9.7 \times 10^4 \text{ m/s}$ .
- 4- The gravitational field intensity of the Earth =  $10 \text{ N/Kg}$ .

**Q5: Write down the slope of the straight line and the mathematical relation for each of the following:**

6)



## Q6: Problems:

- 1- A bicycle rider move in a circular path of radius 40m. With tangential velocity 13.2 m/s. If the force that conserves the circular motion of the bicycle is 377 N. Calculate the mass of the bicycle and the rider.
- 2- A car of mass 905 kg. Moves on a circular path of radius 3.25m. , calculate the centripetal acceleration for the same object if its velocity is doubled and its radius of rotation decreases to its half value.
- 3- An object of mass 2kg is held at the end of a rope and rotate in a horizontal circular path of radius 1.5 m. so , it makes 3 revolutions in one second , calculate :
  - a. The tangential velocity.
  - b. The centripetal acceleration
  - c. The tension in the rope
- 4- Find the mutual attraction between the sun and Jupiter giving that the mass of the sun  $2 \times 10^{30}$ kg , the mass of Jupiter  $1.89 \times 10^{27}$ kg and the distance between their centers is  $7.73 \times 10^{11}$ m (  $G = 6.67 \times 10^{-11}$  N.m<sup>2</sup> /kg<sup>2</sup>)
- 5- A mine is at 500m deep below the surface of Earth. Find the acceleration due gravity inside that mine giving that:  
(G) =  $6.67 \times 10^{-11}$  N.m<sup>2</sup>/kg<sup>2</sup> (R) = 6360km (M) =  $5.98 \times 10^{24}$ kg
- 6- A planet has mass twice that of Earth and a diameter twice that of Earth. Find the ratio of the acceleration due gravity of the two planets.

## Model Answer

### Q1: Complete the following statements:

- 1) Increases – direction
- 2) Decreases – direction
- 3) Direction only
- 4) Its quarter
- 5) Candy floss – rotating barrels in amusement park – Drying cloths.
- 6)  $m \frac{v^2}{r}$
- 7)  $G = \frac{Fr^2}{mM}$
- 8) Equal
- 9) Quarter
- 10)  $N.m^2/kg^2$

### Q2: What's meant by?

- 1) The motion of a body in a circular path with constants speed and changeable direction.
- 2) The force acting continuously in a normal direction of the moving body changing its straight path into circular path.
- 3) The velocity of a body in the tangential direction of circular path at the release moment.

- 4) The force acting normally to the direction of a body moving in a circular path that is inclined to the horizontal towards the center of the curved path helping the body to move in a circular path.
- 5) The acceleration acquired by an object moving in a circular path due to a continuous change in the direction of its velocity.
- 6) The time taken by the body to make one complete revolution.
- 7) A body in the universe attracts another body by a force which is directly proportional to the product of their masses, and inversely proportional to square the distance between them.
- 8) The space in which the gravitational forces appear.
- 9) The gravitational force acting on a mass of 1 kg.
- 10) An object projected at a certain velocity to rotate in a roughly circular path at a constant distance from the Earth's surface.
- 11) The velocity that makes the satellite orbit Earth in a roughly circular path at a constant distance from the Earth's surface.
- 12) They are huge telescopes floating in space in which they can photograph space accurately.

### **Q3: Give Reason:**

- 1) Because when the body moves in a circular path it will acquire a centripetal acceleration which changes the direction of the velocity without changing its magnitude.
- 2) Due to the friction force between the road and car tyers which is normal to the direction of motion of the car causing its motion in a curved path.

- 4) To create a force normal to the direction of motion so the direction of motion changes and the object moves in a circular path.
- 5) Because according to the relation  $F = \frac{mv^2}{r}$  when the car velocity decreases , F also decreases and the car doesn't skid out the road.
- 6) Because the gravitational force is inversely proportional to the square of the distance between the attracted masses.
- 7) Because the orbital velocity can be determined by the relation  $V = \sqrt{\frac{GM}{r}}$   
since M , G are constants then  $V \propto \frac{1}{\sqrt{r}}$
- 8) Because it is used in different fields such as : Communication , photograph space accurately , determine the mineral resources and about the information needed by military and political leaders.

#### Q4: What do we mean when we say.....?

- 1) The force acting in a normal direction on the object's motion to turn it in a circular path is 100N.
- 2) The acceleration acquired by an object due to the change of its velocity direction during its motion in a circular path =  $40\text{m/s}^2$
- 3) The velocity which makes the satellite rotate in a curved path so that its distance from Earth's surface constant =  $9.7 \times 10^4\text{m/s}$
- 4) The attraction force of Earth to an object of mass 1kg = 10N.

## Q5:

a- Slope =  $\frac{v^2}{r} = a$  ( centripetal acceleration )

$$a = \frac{v^2}{r}$$

b- Slope =  $a = \frac{F}{m}$

$$F = ma$$

c- Slope =  $\frac{F}{v^2} = \frac{m}{r}$

$$F = \frac{mv^2}{r}$$

d- Slope =  $\frac{Fr}{m} = V^2$

$$F = \frac{mv^2}{r}$$



## Q6: Problems:

$$1- F = \frac{mv^2}{r}$$

$$m = \frac{Fr}{v^2} = \frac{377 \times 40}{(13.2)^2} = 86.5 \text{ kg.}$$

$$2- V = \sqrt{\frac{Fr}{m}}$$

$$= \sqrt{\frac{2140 \times 3.25 \times 10^3}{905 \times 2\pi}} = 34.97 \text{ m/s}$$

$$3- a- V = \sqrt{\frac{2\pi r}{T}} = \frac{2\pi \times 10^5}{\frac{1}{3}} = 28.26 \text{ m/s}$$

$$b. a = \frac{v^2}{r} = \frac{(28.26)^2}{1.5} = 532.4 \text{ m/s}^2$$

$$c. F = ma = 2 \times 532.4 = 1064.8 \text{ N}$$

$$4- F = G \frac{Mm}{r^2}$$

$$= 6.67 \times 10^{-11} \times \frac{2 \times 10^{30} \times 1.89 \times 10^{27}}{(7.73 \times 10^{11})^2} = 4.22 \times 10^{23} \text{ N}$$

This force keeps Jupiter orbiting the sun.

$$5- g = \frac{GM}{(R-h)^2} = \frac{6.67 \times 10^{-11} \times 5.98 \times 10^{24}}{(6360 \times 10^3 - 500)^2} = 9.86 \text{ N/kg}$$

$$6- M_1 = 2 M_e \quad R_1 = 2 R_e$$

$$\frac{g_1}{g_e} = \frac{M_1 R_e^2}{M_e R_1^2} = \frac{2 M_e R_e^2}{M_e \times 4 R_e^2} = \frac{1}{2}$$

## Part (2)

### Unit (4) Chapter (4) Work and energy

#### Define:

1. Work
  - The dot product of the acting force and the displacement in the direction of the force.
2. Joule
  - The work done by a force of one Newton to move an object through a displacement of one meter in the direction of the force.
3. Energy
  - The ability to do work.
4. Kinetic energy
  - The energy possessed by the object due to its motion.
5. Potential energy
  - The energy stored in the object due to its position or state.

#### Give reasons for:

1. The work done is a scalar quantity although both of force and displacement are vector quantities.
  - Because work is the dot product of force and displacement, which results in scalar.
2. The unit of energy is the same as the unit of work which is joule.
  - Because energy is defined as the ability to do work.
3. Work done can be negative.
  - When the angle  $\theta$  between force direction and the direction of the displacement is  $> 90^\circ$  or the direction of force is opposite to the direction of the displacement.


## Problems:

1. A man of a mass 70 kg ascends the stairs with angle  $60^\circ$  of length 50 m, calculate

the work done ( $g=10 \text{ m/s}^2$ ).

$$F = m g = 70 \times 10 = 700 \text{ N}$$

$$W = Fd \cos \theta = 700 \times 50 \times \cos 30^\circ \\ = 30311 \text{ J.}$$



2. A force affects on a body causing it to displace by 50 m in a direction makes angle  $60^\circ$  with the displacement, if the work done is 2500 j. calculate the force.

$$W = F d \cos \theta$$

$$2500 = F \times 50 \times \cos 60^\circ$$

$$F = 100 \text{ N}$$

3. A force of 5 N acts on a body causing it to displace by 2m, calculate the work done in the following cases:

a. If the acting force is normal to displacement direction.

b. If the acting force makes angle  $30^\circ$  with the direction of displacement.

c. If the acting force is in the same direction of displacement.

$$W = F d \cos \theta$$

a.  $W = 5 \times 2 \times \cos 90^\circ = 0$


b.  $W = 5 \times 2 \times \cos 30^\circ = 8.66 \text{ J}$

c.  $W = 5 \times 2 \times \cos 0^\circ = 10 \text{ J}$

4. A body of mass 50 kg, calculate its height from the Earth's surface if its potential energy at that height equals 2500 j ( $g=10 \text{ m/s}^2$ ).

$$\text{P.E.} = m g h$$

$$h = \text{P.E.} / m g = 2500 / 50 \times 10 = 5 \text{ m}$$



## Chapter (6) Law of conservation of energy

### What is meant by:

1. Law of conservation of energy.

Energy is neither created nor destroyed, but it can be converted from one form to another.

2. Mechanical energy.

The sum of potential energy and kinetic energy of an object.

3. Law of conservation of mechanical energy.

The sum of potential energy and kinetic energy of an object at any point on its path under the effect of gravity only is constant.

### Give reasons for:

1. When an object falls freely its kinetic energy increases.

Because the object's velocity increases due to gravitational acceleration and so,  $K.E. = \frac{1}{2} m V^2$

2. The sum of potential energy and kinetic energy of an object falling freely is constant.

Because according to the conservation law of energy as the kinetic energy increase P.E decrease and vice versa, so their sum stay constant.




### An object is thrown vertically upwards, which of these graphs represents

#### the relation between:

1- P.E. and h ( c )

2- K.E. and h ( a )

3- M.E. and h ( b )



## Problems :

1. An object of mass 0.2 kg is thrown upwards vertically with velocity 20 m/s,  
Neglecting the air resistance, calculate:

- Maximum height the body reaches
- The velocity at a height of 10 m

- a. At the beginning of the movement ( Earth's surface )**

$$M.E. = K.E. + 0 = 1/2 m V^2 = 0.5 \times 0.2 \times 400 = 40 \text{ J}$$

$$\text{At max. height: } M.E. = P.E. + 0 = m g h = 40 \text{ J}$$

$$h = 40 / ( 0.2 \times 10 ) = 20 \text{ m}$$

- b. at height 10 m**

$$P.E. + K.E. = 40 \text{ J}$$

$$0.2 \times 10 \times 10 + 0.5 \times 0.2 \times V^2 = 40 \text{ J} \quad \Longrightarrow \quad V = 14.14 \text{ m/s.}$$

2. An object is thrown upwards vertically with initial velocity 10 m/s, if its potential energy at max. height is 1000 J, find its mass.

$$\text{At max. height: } M.E. = P.E. + 0 = 1000 \text{ J}$$

$$\text{At the Earth's surface: } M.E. = K.E. + 0 \quad \Longrightarrow \quad 1/2 m V^2 = 0.5 \times m \times 100 = 1000 \text{ J}$$


$$m = 20 \text{ kg.}$$

3. A ball is thrown vertically upwards so that its velocity becomes 3 m/s at height 4 m, find the work done to throw the ball if its mass is 0.5 kg and  $g = 10 \text{ m/sec}^2$

$$\text{At height 4 m: } M.E = P.E + K.E.$$

$$M.E. = m g h + 1/2 m V^2 = 0.5 \times 10 \times 4 + 0.5 \times 0.5 \times 9 = 22.25 \text{ J}$$

$$V=0 \quad , \quad K.E. = \text{zero} \quad , \quad P.E. = \text{max.}$$



$$V=20 \quad , \quad K.E. = \text{max.} \quad , \quad P.E. = \text{zero}$$